



Dehissenzen nach Augmentation

Expositionen vermeiden

Ein Beitrag von Dr. Marcus Seiler MSc MSc und Dr. Amely Hartmann

Literaturangabe

- [1] Merli M, Lombardini F, Esposito M. Vertical ridge augmentation with autogenous bone grafts 3 years after loading: resorbable barriers versus titanium-reinforced barriers. A randomized controlled clinical trial. *The International journal of oral & maxillofacial implants.* 2010;25:801-7.
- [2] Lindfors LT, Tervonen EA, Sandor GK, Ylikontiola LP. Guided bone regeneration using a titanium-reinforced ePTFE membrane and particulate autogenous bone: the effect of smoking and membrane exposure. *Oral surgery, oral medicine, oral pathology, oral radiology, and endodontics.* 2010;109:825-30.
- [3] Canullo L, Trisi P, Simion M. Vertical ridge augmentation around implants using e-PTFE titanium-reinforced membrane and deproteinized bovine bone mineral (bio-oss): A case report. *The International journal of periodontics & restorative dentistry.* 2006;26:355-61.
- [4] Boyne PJ, Mikels TE. Restoration of alveolar ridges by intramandibular transposition osseous grafting. *J Oral Surg.* 1968;26:569-76.
- [5] Louis PJ, Gutta R, Said-Al-Naief N, Bartolucci AA. Reconstruction of the maxilla and mandible with particulate bone graft and titanium mesh for implant placement. *Journal of oral and maxillofacial surgery : official journal of the American Association of Oral and Maxillofacial Surgeons.* 2008;66:235-45.
- [6] Rasia-dal Polo M, Poli PP, Rancitelli D, Beretta M, Maiorana C. Alveolar ridge reconstruction with titanium meshes: a systematic review of the literature. *Med Oral Patol Oral Cir Bucal.* 2014;19:e639-46.
- [7] Milinkovic I, Cordaro L. Are there specific indications for the different alveolar bone augmentation procedures for implant placement? A systematic review. *Int J Oral Maxillofac Surg.* 2014;43:606-25.
- [8] Proussaefs P, Lozada J. Use of titanium mesh for staged localized alveolar ridge augmentation: clinical and histologic-histomorphometric evaluation. *J Oral Implantol.* 2006;32:237-47.
- [9] von Arx T, Hardt N, Wallkamm B. The TIME technique: a new method for localized alveolar ridge augmentation prior to placement of dental implants. *The International journal of oral & maxillofacial implants.* 1996;11:387-94.
- [10] Kim YK, Yun PY, Kim SG, Oh DS. In vitro scanning electron microscopic comparison of inner surface of exposed and unexposed nonresorbable membranes. *Oral surgery, oral medicine, oral pathology, oral radiology, and endodontics.* 2009;107:e5-e11.
- [11] Corinaldesi G, Pieri F, Sapigni L, Marchetti C. Evaluation of survival and success rates of dental implants placed at the time of or after alveolar ridge augmentation with an autogenous mandibular bone graft and titanium mesh: a 3- to 8-year retrospective study. *Int J Oral Maxillofac Implants.* 2009;24:1119-28.
- [12] Ciocca L, Mazzoni S, Fantini M, Persiani F, Baldissara P, Marchetti C et al. A CAD/CAM-prototyped anatomical condylar prosthesis connected to a custom-made bone plate to support a fibula free flap. *Med Biol Eng Comput.* 2012;50:743-9.
- [13] Wang G, Li J, Khadka A, Hsu Y, Li W, Hu J. CAD/CAM and rapid prototyped titanium for reconstruction of ramus defect and condylar fracture caused by mandibular reduction. *Oral surgery, oral medicine, oral pathology and oral radiology.* 2012;113:356-61.
- [14] Seiler M, KPW, Peetz M., Hartmann A.G. Customized Lattice Structure in Reconstruction of Three-Dimensional Alveolar Defects: A Technical Note. *International Journal of Computerized Dentistry.* 2018;in Press.
- [15] Seiler M, Kammerer PW, Peetz M, Hartmann AG. Customized Titanium Lattice Structure in Three-Dimensional Alveolar Defect: An Initial Case Letter. *The Journal of oral implantology.* 2018;44:219-24.
- [16] Seiler M PM, Hartmann A, Witkowski R. Individualized CAD/CAM-produced titanium scaffolds for alveolar bone augmentation: a retrospective analysis of dehiscence events in relation to demographic and surgical parameters. *J Oral Science Rehabilitation.* 2018;4(1):38-46.
- [17] Ciocca L, Fantini M, De Crescenzo F, Corinaldesi G, Scotti R. Direct metal laser sintering (DMLS) of a customized titanium mesh for prosthetically guided bone regeneration of atrophic maxillary arches. *Medical & biological engineering & computing.* 2011;49:1347-52.
- [18] Jung GU, Jeon JY, Hwang KG, Park CJ. Preliminary evaluation of a three-dimensional, customized, and preformed titanium mesh in peri-implant alveolar bone regeneration. *J Korean Assoc Oral Maxillofac Surg.* 2014;40:181-7.
- [19] Leghissa GC, Zaffe D, Assenza B, Botticelli AR. Guided bone regeneration using titanium grids: report of 10 cases. *Clin Oral Implants Res.* 1999;10:62-8.
- [20] Degidi M, Scarano A, Piattelli A. Regeneration of the alveolar crest using titanium micromesh with autologous bone and a resorbable membrane. *J Oral Implantol.* 2003;29:86-90.
- [21] Singare S, Dichen L, Bingheng L, Yanpu L, Zhenyu G, Yaxiong L. Design and fabrication of custom mandible titanium tray based on rapid prototyping. *Medical engineering & physics.* 2004;26:671-6.
- [22] Hartmann A, Seiler M. Positive Effects of A- and I-PRF in management of titanium mesh exposures. 2019;submitted.

