



Präparationsform zur minimalinvasiven Versorgung der Unterkiefer-Frontzähne

Der Wellenschliff

Ein Beitrag von Dr. Michael Fischer und Ztm. Benjamin Votteler

Literaturangabe

- [1] Ahlers MO, Möller K: Repositions- Onlays und Veneers zur atraumatischen Restauration einer physiologischen Kiefer- und Kondylenposition. Quintessenz 62, 211-222, 2011
- [2] Ahlers MO, Vahle-Hinz K, Rybczynski AM, Jakstat HA. Semipermanente und permanente Übertragung der Schienenposition mittels Repositions-Onlays und -Veneers: Varianten und Überlebensdauer. Bad Homburg: 44. Jahrestagung der Deutschen Gesellschaft für Funktionsdiagnostik und -therapie (DGFDT), 2011
- [3] Arnetzl, G.V., Arnetzl, G. 2006. Design of preparations for all-ceramic inlay materials. International journal of computerized dentistry. 9(4):289-298
- [4] Burke FJ. Survival rates for porcelain laminate veneers with special reference to the effect of preparation in dentin: a literature review. J Esthet Restor Dent 2012;24:257-265
- [5] Carvalho RM, Santiago SL, Fernandes CA, Suh BI, Pashley DH (2000). Effects of prism orientation on tensile strength of enamel. J Adhes Dent 2(4):251-257
- [6] Clausen JO, Abou Tara M, Kern M. Dynamic fatigue and fracture resistance of non-retentive all-ceramic full-coverage molar restorations. Influence of ceramic material and preparation design. Dent Mater 2010; 26:533-538
- [7] Edelhoff D, Beuer F, Stimmelmayer M, Güth J.F., Minimalinvasive Behandlungskonzepte in der festsitzenden Prothetik, Quintessenz 2014;65(5):589-600
- [8] Edelhoff D, Sorensen JA.: Tooth structure removal associated with various preparation designs for posterior teeth. Int J Periodontics Restorative Dent. 2002 Jun;22(3):241-9
- [9] Frankenberger R, Tay FR (2005). Self-etch vs etch-and-rinse adhesives: effect of thermo-mechanical fatigue loading on marginal quality of bonded resin composite restorations. Dent Mater 21(5):397-412
- [10] Garcia-Godoy F, Krämer N, Feilzer AJ, Frankenberger R (2010). Long-term degradation of enamel and dentin bonds: 6-year results in vitro vs. in vivo. Dent Mater 26(11):1113-1118.
- [11] Gurel G, Sesma N, Calamita MA, Coachman C, Morimoto S. Influence of enamel preservation on failure rates of porcelain laminate veneers. Int J Periodontics Restorative Dent 2013;33:31-39
- [12] Hajto J, Frontzahnrestaurationen: Sind Veneers die beste Wahl? Quintessenz 2010;61(5):521-528
- [13] Körber K: Zahnärztliche Prothetik. Thieme Verlag, Stuttgart (1995)
- [14] Magne J Douglas WH. Design optimization and evolution of bonded ceramics for the anterior dentition: a finite-element analysis. Quintessenz Int 1999;30:661-672
- [15] Schulte W: Die exzentrische Okklusion. Quintessenz Verlags-GmbH, Berlin (1983)
- [16] Strub JR, Türp JC, Witkowski S, Hürzeler MB, Kern M. Curriculum Prothetik. Band I. 2.Aufl. Berlin: Quintessenz Verlags-GmbH; 1999
- [17] Van Meerbeek B, De Munck J, Yoshida Y, Inoue S, Vargas M, Vijay P et al. (2003b). Buonocore memorial lecture. Adhesion to enamel and dentin: current status and future challenges. Oper Dent 28(3):215-235

