

## Treatment approach for the severely atrophied maxilla

# Use of platelet-rich fibrin in combination with a bovine bone substitute material

PROFESSOR SHAHRAM GHANAATI, DR SARAH AL-MAAWI, DR TORSTEN CONRAD, ALL FRANKFURT AM MAIN, GERMANY

- [1] R. Pistilli, L. Signorini, A. Pisacane, G. Lizio, and P. Felice, "Case of severe bone atrophy of the posterior maxilla rehabilitated with blocks of equine origin bone: histological results.," *Implant Dent.*, vol. 22, no. 1, pp. 8–15, Feb. 2013.
- [2] J. Lorenz et al., "Do clinical and radiological assessments contribute to the understanding of biomaterials? Results from a prospective randomized sinus augmentation split-mouth trial," *J. Oral Implantol.*, vol. 44, no. 1, 2018.
- [3] S. Ghanaati, M. Barbeck, P. Booms, J. Lorenz, C. J. Kirkpatrick, and R. A. Sader, "Potential lack of 'standardized' processing techniques for production of allogeneic and xenogeneic bone blocks for application in humans," *Acta Biomater.*, vol. 10, no. 8, pp. 3557–3562, 2014.
- [4] J. Lorenz et al., "TRAP-Positive Multinucleated Giant Cells Are Foreign Body Giant Cells Rather Than Osteoclasts: Results From a Split-Mouth Study in Humans," *J. Oral Implantol.*, vol. 41, no. 6, pp. e257–e266, 2015.
- [5] C. M. Younger EM1, "Morbidity at bone graft donor sites. - PubMed - NCBI." [Online]. Available: <https://www.ncbi.nlm.nih.gov/pubmed/?term=10.%09Younger+EM%2C+Chapman+MW%3A+Morbidity+at+bone+graft+donor+sites.+J+Orthop+Trauma+1989%3B+3%3A192-5>.
- [6] E. Dohle, K. El Bagdadi, R. Sader, J. Choukroun, C. James Kirkpatrick, and S. Ghanaati, "PRF-based matrices to improve angiogenesis in an in vitro co-culture model for bone tissue engineering," *J. Tissue Eng. Regen. Med.*, 2017.
- [7] K. El Bagdadi et al., "Reduction of relative centrifugal forces increases growth factor release within solid platelet-rich-fibrin (PRF)-based matrices: a proof of concept of LSCC (low speed centrifugation concept)," *Eur. J. Trauma Emerg. Surg.*, Mar. 2017.
- [8] S. Wend et al., "Reduction of the relative centrifugal force influences cell number and growth factor release within injectable PRF-based matrices," *J. Mater. Sci. Mater. Med.*, vol. 28, no. 12, p. 188, Dec. 2017.
- [9] E. Kobayashi et al., "Comparative release of growth factors from PRP, PRF, and advanced-PRF," *Clin. Oral Investig.*, pp. 1–8, 2016.
- [10] R. Verboket et al., "Influence of concentration and preparation of platelet rich fibrin on human bone marrow mononuclear cells (in vitro)," *Platelets*, pp. 1–10, Oct. 2018.
- [11] J. Choukroun and S. Ghanaati, "Reduction of relative centrifugation force within injectable platelet-rich-fibrin (PRF) concentrates advances patients' own inflammatory cells, platelets and growth factors: the first introduction to the low speed centrifugation concept," *Eur. J. Trauma Emerg. Surg.*, vol. 44, no. 1, pp. 87–95, Feb. 2018.
- [12] Al-Maawi et al., "Homogeneous pressure influences the growth factor release profiles in solid platelet-rich fibrin matrices and enhances vascular endothelial growth factor release in the solid platelet-rich fibrin plugs," *Int. J. Growth Factors Stem Cells Dent.*, vol. 1, no. 1, p. 8, 2018.
- [13] C. W. Digiovanni and J. M. Petricek, "The Evolution of rhPDGF-BB in Musculoskeletal Repair and its Role in Foot and Ankle Fusion Surgery," *Foot Ankle Clin. NA*, vol. 15, pp. 621–640, 2010.
- [14] E. Dohle, K. El Bagdadi, R. Sader, J. Choukroun, C. James Kirkpatrick, and S. Ghanaati, "PRF-based matrices to improve angiogenesis in an in vitro co-culture model for bone tissue engineering," *J. Tissue Eng. Regen. Med.*, May 2017.
- [15] S. Wend et al., "Reduction of the relative centrifugal force influences cell number and growth factor release within injectable PRF-based matrices," *J. Mater. Sci. Mater. Med.*, vol. 28, no. 12, 2017.

- [16] A. Acocella, R. Bertolai, J. Nissan, E. Ellis, and R. Sacco, "Maxillary sinus lift using fresh frozen bone chips in presence of sinus cyst: clinical and histological report.," *Cell Tissue Bank.*, vol. 13, no. 2, pp. 327–32, Jun. 2012.
- [17] S. Ghanaati et al., "Synthetic bone substitute material comparable with xenogeneic material for bone tissue regeneration in oral cancer patients: First and preliminary histological, histomorphometrical and clinical results.," *Ann. Maxillofac. Surg.*, vol. 3, no. 2, pp. 126–38, 2013.
- [18] N. Hatano, Y. Shimizu, and K. Ooya, "A clinical long-term radiographic evaluation of graft height changes after maxillary sinus floor augmentation with a 2 : 1 autogenous bone/xenograft mixture and simultaneous placement of dental implants," *Clin. Oral Implants Res.*, vol. 15, no. 3, pp. 339–345, Jun. 2004.
- [19] G. M. Raghoobar, P. Onclin, G. C. Boven, A. Vissink, and H. J. A. Meijer, "Long-term effectiveness of maxillary sinus floor augmentation A systematic review and meta-analysis," *J. Clin. Periodontol.*, Jan. 2019.
- [20] K. Sagheb, E. Schiegnitz, M. Moergel, C. Walter, B. Al-Nawas, and W. Wagner, "Clinical outcome of alveolar ridge augmentation with individualized CAD-CAM-produced titanium mesh.," *Int. J. Implant Dent.*, vol. 3, no. 1, p. 36, Dec. 2017.
- [21] J. Lorenz et al., "Allogeneic bone block for challenging augmentation—a clinical, histological, and histomorphometrical investigation of tissue reaction and new bone formation," *Clin. Oral Investig.*, 2018.
- [22] T. Asaka, N. Ohga, Y. Yamazaki, J. Sato, C. Satoh, and Y. Kitagawa, "Platelet-rich fibrin may reduce the risk of delayed recovery in tooth-extracted patients undergoing oral bisphosphonate therapy: a trial study," 2004.
- [23] S. Ghanaati et al., "Fifteen years of platelet rich fibrin (PRF) in dentistry and oromaxillofacial surgery: How high is the level of scientific evidence?," *J. Oral Implantol.*, p. aaid-joi-D-17-00179, Jun. 2018.
- [24] A. Simonpieri, J. Choukroun, M. Del Corso, G. Sammartino, and D. M. Dohan Ehrenfest, "Simultaneous sinus-lift and implantation using microthreaded implants and leukocyte- and platelet-rich fibrin as sole grafting material: a six-year experience.," *Implant Dent.*, vol. 20, no. 1, pp. 2–12, Feb. 2011.